

ON THE SEVERI VARIETIES OF SURFACES IN $\mathbf{P}^3$

L.CHIANTINI

Universita' di Siena - Dipartimento di Matematica
Via del Capitano, 15 53100 SIENA (Italy)
e-mail: chiantini@unisi.it

C.CILIBERTO

Dipartimento di Matematica - Universita' di Roma - Tor Vergata
loc. La Romanina, 00133 ROMA (Italy)
e-mail: ciliberto@axp.mat.utorrm.it

ABSTRACT. For a smooth surface S in $\mathbf{P}^3$ of degree d and for positive integers n, δ , the Severi variety $V_{n,\delta}^0(S)$ is the subvariety of the linear system $|O_S(n)|$ which parametrizes curves with δ nodes. We show that for S general, $n \geq d$ and for all δ with $0 \leq \delta \leq \dim(|O_S(n)|)$, then $V_{n,\delta}^0(S)$ has at least one component which is reduced, of the expected dimension $\dim(|O_S(n)|) - \delta$. We also construct examples of reducible Severi varieties on general surfaces of degree $d \geq 8$.

INTRODUCTION

Nodal curves in the plane have been classically extensively studied. The variety $V_{n,\delta}$ of irreducible plane curves of degree n with δ nodes is called the *Severi variety* of type (n, δ) . Severi claimed in [Se] the irreducibility of $V_{n,\delta}$, but his proof was not complete. This result has been more recently established by J. Harris in [H].

It is natural to extend the notion of Severi varieties to curves on any smooth projective surface S . Take an *effective* line bundle $\mathcal{L}$ on S , i.e. a line bundle such that the linear system $|\mathcal{L}|$ is not empty. Let $p_{\mathcal{L}}$ be the arithmetic genus of a curve $C \in |\mathcal{L}|$ and let δ be an integer such that $0 \leq \delta \leq p_{\mathcal{L}}$. It makes sense to consider the subscheme $V_{\mathcal{L},\delta}^0$ of the projective space $|\mathcal{L}|$ consisting of all curves $C \in |\mathcal{L}|$ which are irreducible and have δ nodes and no other singular point. This scheme is locally closed in $|\mathcal{L}|$, and we can consider its closure $V_{\mathcal{L},\delta}$ in $|\mathcal{L}|$, which we call the *Severi variety* of curves with δ nodes in $|\mathcal{L}|$. Often we will also refer to the open dense subset $V_{\mathcal{L},\delta}^0$ as to the Severi variety.

As for the plane case, one may ask several questions concerning these Severi varieties. For example:

- (i) *the existence problem*: when is $V_{\mathcal{L},\delta}$ not empty?
- (ii) *the dimension problem*: what are the dimensions of the components of $V_{\mathcal{L},\delta}$?
- (iii) *the irreducibility problem*: when is $V_{\mathcal{L},\delta}$ irreducible?
- (iv) *the enumeration problem*: what is the degree of $V_{\mathcal{L},\delta}$ as a variety in $|\mathcal{L}|$?

There is a natural approach to the dimension problem which uses infinitesimal deformations of nodal curves. Let C be a curve in $V_{\mathcal{L},\delta}^0$, which we thus suppose not

empty, and let Z be its set of nodes. One knows (see [S]) that the Zariski tangent space to $V_{\mathcal{L},\delta}^0$ at C is

$$T_{V_{\mathcal{L},\delta}^0,C} \simeq H^0(S, \mathcal{I}_{Z,S} \otimes \mathcal{L})/(C)$$

and the corresponding obstruction space is $H^1(S, \mathcal{I}_Z \otimes \mathcal{L})$. Let us consider the number $c_{Z,\mathcal{L}}$ of independent conditions which Z imposes to the linear system $|\mathcal{L}|$, i.e. the rank of the natural restriction map

$$H^0(S, \mathcal{L}) \rightarrow H^0(Z, \mathcal{O}_Z)$$

Then one has

$$\begin{aligned} \dim_C V_{\mathcal{L},\delta}^0 &\leq \dim T_{V_{\mathcal{L},\delta}^0,C} = \\ &= h^0(S, \mathcal{I}_Z \otimes \mathcal{L}) - 1 = \dim |\mathcal{L}| - c_{Z,\mathcal{L}} \end{aligned}$$

On the other hand one has also

$$\begin{aligned} \dim_C V_{\mathcal{L},\delta}^0 &\geq h^0(S, \mathcal{I}_Z \otimes \mathcal{L}) - 1 - h^1(S, \mathcal{I}_Z \otimes \mathcal{L}) - 1 = \\ &= \dim |\mathcal{L}| - \delta \end{aligned}$$

and $\max\{\dim |\mathcal{L}| - \delta, -1\}$ is the so-called *expected dimension* of $V_{\mathcal{L},\delta}^0$ at C . In conclusion $V_{\mathcal{L},\delta}^0$ is smooth of codimension δ in $|\mathcal{L}|$ at C if and only if $\delta \leq \dim |\mathcal{L}|$ and Z imposes independent conditions to $|\mathcal{L}|$. In this case one says that $V_{\mathcal{L},\delta}^0$ and $V_{\mathcal{L},\delta}$ are *regular* at C . If instead $\dim_C V_{\mathcal{L},\delta}^0 > \dim |\mathcal{L}| - \delta$ we will say that $V_{\mathcal{L},\delta}^0$ is *superabundant* at C .

Regularity is a very strong property. In fact, again by using infinitesimal deformation arguments (see [S]), one sees that it implies that *the nodes of C can be independently smoothed*. In other words for any subset Z' of Z , of order δ' we can find a family of curves in $|\mathcal{L}|$, parametrized by a disc Δ , whose general member C_t , $t \in \Delta - \{0\}$, is nodal with δ' nodes, its special fibre over 0 is C , and the set of nodes of the general member specializes to Z' when C_t specializes to C .

It is easy to see that if S is a rational surface and $C \cdot K_S < 0$ for $C \in |\mathcal{L}|$, then, if C sits in $V_{\mathcal{L},\delta}^0$, its set of nodes Z imposes independent conditions to $|\mathcal{L}|$, hence all components of $V_{\mathcal{L},\delta}^0$ are regular. This applies in particular to the plane or, more generally, to the del Pezzo surfaces, for which one has that all components of the Severi varieties are regular.

A similar argument also works for other regular surfaces like K3 or Enriques surfaces, for which again one proves that all components of the Severi varieties are regular (see [AC], [T1,2]).

Let us now turn to the existence problem. In the plane case one takes advantage from the fact that there are rational nodal curves of any degree n , namely the general projections to a plane of rational normal curves of degree n in $\mathbf{P}^n$. By smoothing some of their nodes one proves the non emptiness of $V_{n,\delta}^0$ for all $\delta \leq \frac{(n-1)(n-2)}{2}$.

An investigation about non emptiness of $V_{\mathcal{L},\delta}^0$ for del Pezzo surfaces is contained in the recent paper [GLS].

The existence problem becomes more difficult for other surfaces, like K3 or Enriques surfaces. For example take an effective line bundle $\mathcal{L}$ on a K3 surface S . Notice that $p_{\mathcal{L}} = \dim|\mathcal{L}|$, hence we may expect the existence of rational nodal curves in $|\mathcal{L}|$. They would be clearly finitely many and regular, and this in turn would imply that $V_{n,\delta}^0$ is regular for all integers δ such that $0 \leq \delta \leq \dim|\mathcal{L}|$.

Recently Xi Chen proved in [Ch] the following result: *for all $g \geq 3$, on the general K3 surface S of the principal series in $\mathbf{P}^g$ there are rational nodal curves in the linear system $|\mathcal{O}_S(n)|$ for all $n > 0$.* We will comment on this theorem in §2, where we will recall the main idea of its proof in the case $g = 3$.

In recent times there has also been a lot of interest about the enumerative problem. We recall that the degree of the Severi variety of rational curves of degree n has been computed by Kontsevich in [K], whereas Caporaso and Harris gave in [CH] a general formula for the degree of all Severi varieties in the plane. These formulas were in principle also contained in the paper [R] by Ziv Ran.

We also recall the paper [YZ] by Yau and Zaslow, in which one finds a suggestion for the computation of the number of rational hyperplane sections of the general K3 surface S of the principal series in $\mathbf{P}^g$. For an algebro-geometric account of [YZ] one can look at Beauville's paper [B]. Unfortunately there are still some missing details in order to make Yau and Zaslow's beautiful argument work: among these one should prove that *all* rational hyperplane sections of the general K3 surface S of the principal series in $\mathbf{P}^g$ are nodal. This has been announced, but still not completely proved, by Chen in [Ch].

In the present paper we consider Severi varieties for surfaces of degree d in $\mathbf{P}^3$ and in particular we address the existence and dimension problems. Since the cases $d \leq 4$ are covered by the aforementioned results, we consider the case $d \geq 5$, which is interesting since the surfaces in question are of general type, and for such surfaces very little is known about the structure of their Severi varieties.

Let S be a smooth surface of degree d in $\mathbf{P}^3$. We use the notation $V_{n,\delta}(S)$ and $V_{n,\delta}^0(S)$ to denote the Severi varieties of curves in $|\mathcal{O}_S(n)|$. A recent result by the first author and Sernesi [CS] says that *all* components of $V_{n,\delta}^0(S)$ are regular if δ is *small* with respect to d and n , for instance if $\delta < \frac{nd(n-2d+8)}{4}$.

On the other hand, if we are on a surface S of degree d , we could in principle expect the existence of regular components for all δ such that $0 \leq \delta \leq \dim|\mathcal{O}_S(n)| := N(d, n)$. This is exactly what we investigate in the present paper.

Our results are of two types. First, in §1, we give examples of superabundant components $V_{n,\delta}^0(S)$ on any surface S of degree d . Our examples are of two kinds. One kind consists of not empty components of $V_{n,\delta}^0(S)$ with $\delta > N(d, n)$, which are clearly superabundant. The second kind instead consists of superabundant components in the range $\delta \leq N(d, n)$, $n \geq d$. Both examples are obtained by intersecting cones of a suitable family with S .

After having recalled in §2 Chen's theorem mentioned above, in §3 we extend it to general surfaces of higher degrees. Our main result is theorem 3.1, which says that if S is a general surface of degree $d \geq 5$ in $\mathbf{P}^3$, for any $n > 0$ and for any δ such that $0 \leq \delta \leq N(d, n)$ there exist regular components of $V_{n,\delta}^0(S)$. The proof is by induction, the first step of which is Chen's theorem. Then we use degeneration and we prove the existence of surfaces of degree d which split into a general surface of degree $d-1$ plus a general tangent plane of it. We can prove that on such reducible surfaces we have *limit nodal curves* with good properties. By partially smoothing

these curves we find our regular components of the Severi varieties on a general surface.

We finish by noticing that, putting together our main theorem 3.1 and the examples of §1, we conclude with the surprising result that on a general surface S of large enough degree in $\mathbf{P}^3$ some Severi varieties are reducible.

We pose however the following question: *is it true that for S general of degree d and for all δ such that $0 \leq \delta \leq N(d, n)$, there is only one regular irreducible component of $V_{n, \delta}^0(S)$?*

We also have a question about superabundant components, which naturally arises by looking at our examples in §2: *is it true that for S general all the superabundant components of $V_{n, \delta}^0(S)$ are such that their general curve is cut out on S by a surface which is singular in codimension one?*

In conclusion, the second author would like to thank J. Harris for several interesting discussions on the subject of this paper. Both authors wish to thank the referee for useful remarks on the first version of the paper.

0. NOTATION

We work over the complex field $\mathbf{C}$. In this paper we will use the following notation:

S will be a general surface of degree d in $\mathbf{P}^3$.

$\mathbf{P}_d = |\mathcal{O}_{\mathbf{P}^3}(d)|$ is the linear system of surfaces of degree d in $\mathbf{P}^3$. We have:

$$\dim \mathbf{P}_d = \frac{d^3 + 6d^2 + 11d}{6}.$$

$p_a(d, n) = \frac{dn}{2}(n + d - 4) + 1$ is the arithmetic genus of the divisors in $|\mathcal{O}_S(n)|$.

$V_{n, \delta}(S)$ is the Severi variety of curves in $|\mathcal{O}_S(n)|$ with δ nodes, i.e. the closure in $|\mathcal{O}_S(n)|$ of the scheme $V_{n, \delta}^0(S)$ of irreducible curves with δ distinct nodes and no other singularity. Notice that $V_{n, \delta}^0(S)$ is open in $V_{n, \delta}(S)$. If there is no danger of confusion, we will write $V_{n, \delta}$ and $V_{n, \delta}^0$ instead of $V_{n, \delta}(S)$ and $V_{n, \delta}^0(S)$.

$N(d, n) = h^0 \mathcal{O}_S(n) - 1 = \dim |\mathcal{O}_S(n)|$. We have:

$$N(d, n) = \begin{cases} \frac{nd}{2}(n - d + 4) + \frac{d^3 - 6d^2 + 11d}{6} - 1 & \text{for } n \geq d \\ \dim \mathbf{P}_n & \text{for } n < d \end{cases}$$

We say that $C \subset S$ is a *nodal curve* if it is irreducible, with only nodes as singularities. $V_{n, \delta}^0$ parametrizes all nodal curves of geometric genus $p_a(n, d) - \delta$ in the system $|\mathcal{O}_S(n)|$.

Notice that on some dense Zariski open subset U of $\mathbf{P}_d$ we have a universal surface $\mathcal{S} \rightarrow U$ and a universal linear system $|\mathcal{O}_{\mathcal{S}}(n)|$, and one can also define the universal algebraic schemes $\mathcal{V}_{n, \delta}$ and $\mathcal{V}_{n, \delta}^0$ corresponding to the Severi varieties.

1. SUPERABUNDANT COMPONENTS OF SEVERI VARIETIES

As we said in the introduction, in the plane, i.e. when $d = 1$, it is classically known that the Severi varieties $V_{n, \delta}$, when not empty, all have the expected dimension $N(d, n) - \delta$. This result extends to every Del Pezzo surface. In particular it applies also to quadrics and cubics, i.e. to the cases $d = 2, 3$.

On the other hand, when the degree d increases and S is of general type, the situation is no longer so simple. On a general surface of degree $d \gg 0$ the Severi varieties may have several components, with different dimensions.

In particular, the following examples show the existence of superabundant components of $V_{n,\delta}^0$. Later on we will see that for the values of n , d and δ considered in example (1.2) below there are also regular components, hence the corresponding Severi varieties turn out to be reducible.

Example 1.1. Fix $d \geq 20$ and let S be a smooth surface of degree d in $\mathbf{P}^3$. Consider $n = 3$, $\delta = 20$. We have $N(d, 3) = 19$ so the expected dimension of $V_{n,\delta}^0$ is negative.

On the other hand, $V_{n,\delta}^0$ is non empty: indeed, take the intersection of S with a general cone over a singular plane cubic curve.

Even when the expected dimension is positive, it may happen that $V_{n,\delta}^0$ has superabundant components. One can produce such examples using again cones.

Example 1.2. Take a smooth surface S of degree $d \geq 8$ and take $n \gg d$, n even.

On a general plane there is a family of dimension $(nd - n - 2)/2$ of irreducible nodal curves of degree n having exactly

$$\alpha = \frac{n^2 - 3n + 2}{2} - \frac{nd}{2} + \frac{7n}{2}$$

nodes. Fixing a general $P \in \mathbf{P}^3$, we have a family of cones with vertex at P over these plane curves. Such cones will have α double lines. Intersecting these cones with S , we get a family contained in $V_{n,\delta}^0$ with $\delta = d\alpha$, whose dimension is at least $(nd - n - 2)/2$. In fact two cones cannot intersect S in the same curve C , since C and P determine the cone.

One computes the expected dimension of $V_{n,d\alpha}^0$ which is:

$$N(d, n) - d\alpha = \frac{d^3 - 12d^2 + 11d - 6}{6}$$

Notice that it becomes soon smaller than $(nd - n - 2)/2$ when n is large with respect to d .

Remember from the Introduction that there are no superabundant components of Severi varieties on a quartic surface. In particular, for $\delta = N(4, n)$, then $V_{n,\delta}^0$ is a finite set of points. There is a general, easy criterion for the finiteness of $V_{n,\delta}^0$, when δ is large.

Proposition 1.4. *If $2\delta > n^2d$ then $V_{n,\delta}^0$ is formed by isolated points.*

Proof. Assume that $V_{n,\delta}^0$ has a component of positive dimension. This implies the existence of a non trivial family of nodal curves, with δ nodes, in $|\mathcal{O}_S(n)|$. If C is a general element of this family and Z is its set of nodes, then there exists a non-zero element in $T_{C, V_{n,\delta}^0} \simeq H^0 \mathcal{I}_{Z,S}(n)/(C)$, i.e. there exists another curve $C' \in |\mathcal{O}_S(n)|$ passing through the nodes of C . Then one has:

$$2\delta \leq C \cdot C' = n^2d.$$

2. NODAL CURVES ON GENERAL QUARTIC SURFACES

In this paragraph we consider the case $d = 4$. Notice that $N(4, n) = 2n^2 + 1 = p_a(4, n)$ and, if $N(4, n) - \delta \geq 0$, then every component of $V_{n, \delta}^0$ is regular.

In the case S is general, the existence problem, i.e. the non-emptiness of $V_{n, \delta}^0$, follows from a general theorem by X. Chen [Ch], whose result, in the present situation, boils down to the following:

Theorem 2.1. [Ch] *Let S be a general quartic surface in $\mathbf{P}^3$. Then for all $n > 0$, the linear system $|\mathcal{O}_S(n)|$ contains irreducible nodal rational curves.*

There are two consequences of Chen's theorem 2.1:

- (i) $V_{n, \delta}^0 \neq \emptyset$ for all $\delta \leq N(4, n)$ (see the Introduction);
- (ii) looking at the universal Severi varieties, the theorem implies that $\mathcal{V}_{n, \delta}^0$, for $\delta = N(4, n)$, has a component which dominates $\mathbf{P}_4$, with 0-dimensional general fiber.

It is necessary for our purposes to briefly recall the idea of Chen's proof.

Let $S_0 = Q_1 \cup Q_2$ be a special quartic in $\mathbf{P}^3$ formed by the union of two general quadrics Q_i , $i = 1, 2$, being general, is smooth, therefore it has two rulings of lines, which we denote by H_j^i , $j = 1, 2$. Moreover the intersection E of Q_1 and Q_2 is a smooth elliptic quartic curve in $\mathbf{P}^3$.

We consider on S_0 the curve C_0 defined as follows:

$$C_0 = C_0^1 \cup C_0^2$$

with C_0^i sitting on Q_i , $i = 1, 2$, defined as:

$$C_0^i = C_1^i \cup \dots \cup C_n^i$$

where $C_j^i \in |H_1^i|$, $j = 1, \dots, n-1$, are distinct lines and $C_n^i \in |H_1^i + nH_2^i|$ is a smooth irreducible curve, subject to the following conditions:

- (i) the line C_j^1 , $j = 1, \dots, n-1$, intersects E at two distinct points $q_{2(j-1)}, q_{2j-1}$;
- (ii) the line C_j^2 , $j = 1, \dots, n-1$, intersects E at the two distinct points q_{2j-1}, q_{2j} ;
- (iii) the curve C_n^i passes through $q_{2(n-1)}$ and has a contact of order $2n+1$ with E at point r different from all the points q_h , $h = 0, \dots, 2(n-1)$;
- (iv) the curve C_n^2 passes through q_0 and has a contact of order $2n+1$ with E at r .

Notice that the linear systems $|H_1^i + nH_2^i|$, $i = 1, 2$, cut on E a complete linear series of degree $2n+2$ and dimension $2n+1$. Taking this into account, after one moment of reflection the reader will see that such a configuration exists, and that C_0 is a divisor in $|\mathcal{O}_{S_0}(n)|$.

We will refer to C_0 as to a *Chen's curve*.

Now one takes a general quartic surface S passing through q_0 , and therefore cutting E in fifteen more distinct points $p_1, \dots, p_{15}$ different from the points q_h , $h = 1, \dots, 2(n-1)$, and r . One takes the pencil generated by S_0 and S , which we briefly denote by $S_t = tS + S_0$. General facts about deformations of singular curves (see [Ch], [R]), ensure that, for t in a small disc Δ around 0, C_0 deforms to an irreducible nodal curve C_t on S_t in the linear system $|\mathcal{O}_{S_t}(n)|$, having $2n^2 + 1$ distinct nodes, and therefore being rational. The limits of the nodes of C_t , when t tends to 0 are:

- (i) the $n(n-1)$ nodes of C_0^1 arising from the intersections of $C_1^1, \dots, C_{n-1}^1$ with C_n^1 ;
- (ii) the $n(n-1)$ nodes of C_0^2 arising from the intersections of $C_1^2, \dots, C_{n-1}^2$ with C_n^2 ;
- (iii) the point q_0 ;
- (iv) the $2n+1$ -tacnode r where $2n$ nodes of C_t coalesce.

Let now S be a smooth surface in $\mathbf{P}^3$ and let us take a curve C of degree m on S . Let π be the general tangent plane to S . Notice that π is not tangent to C , since the dual variety V^* of a given projective variety V determines V . Then the divisor Γ cut out by π on C is formed by m distinct points. Let U be the not empty open subset U of S^* such that for all $\pi \in U$ this happens. We can consider the incidence correspondence

$$X = \{(x, \pi) \in C \times U : x \in \pi\}$$

The projection $p : X \rightarrow U$ is a covering of degree m , whose monodromy group, which is a subgroup of the full symmetric group S_m on m objects, we denote by $G(C, S)$.

The following result will be an important ingredient for our construction of regular components of Severi varieties with the maximum number $N(d, n)$ of nodes.

Theorem 2.2. *If S is a general quartic surface, for all n and $\delta \leq N(4, n)$ there is an irreducible component of $V_{n, \delta}^0(S)$ such that for C general in this component one has $G(C, S) = S_{4n}$.*

Proof. Since any irreducible nodal rational curve on S is the limit of nodal curves with $\delta \leq N(4, n)$ nodes, it clearly suffices to prove that on a general quartic surface S there is some rational nodal curve C in $|\mathcal{O}_S(n)|$ for which $G(C, S) = S_{4n}$.

We consider the setting of Chen's proof of theorem (2.1) and we keep the notation introduced above. We will prove that a rational curve C_t on S_t , $t \in \Delta - \{0\}$, which tends to a Chen's curve, is such that $G := G(C_t, S_t)$, i.e. the monodromy group of the corresponding cover $p_t : X_t \rightarrow U_t$, is the full symmetric group S_{4n} . In order to do so, we will prove that G contains all transpositions.

First we prove that G contains a simple transposition. To prove this, it suffices to prove that the monodromy group $G_0 = G(C_0, S_0)$ of the limit covering of $p_t : X_t \rightarrow U_t$ contains a simple transposition. When S_t tends to S_0 , the dual variety S_t^* breaks up in several components, among which there are the dual planes corresponding to the points p_i , $i = 1, \dots, 15$. Actually we may assume that p_1 is a general point of E . Then the general tangent plane σ to C_n^2 through p_1 is simply tangent to C_n^2 at some point w and meets transversally the curve C_0 at the remaining intersection points. The existence of such a simply tangent plane yields a transposition in G_0 as proved in [ACGH], pg. 112. Indeed a plane π_0 passing through p_1 and sufficiently close to σ cuts C_0 transversally, and two points of the intersection x_0, y_0 come together at w when π_0 tends to σ . These two points are exchanged by a simple trasposition in G_0 .

Let us prove now that G contains all transpositions. Let us take C_t close to C_0 and π_t tangent to S_t close to π_0 . Let us take a_t, b_t two points in $\Gamma_t = \pi_t \cap C_t$. Suppose that the limits of these two points, when t tends to 0, are two points a_0, b_0 of C_n^2 . Then we easily find a permutation in G_0 which sends the pair (a_0, b_0) to the pair (x_0, y_0) , simply by moving π_0 inside the dual plane to p_1 , and by letting a_0 tend to x_0 and b_0 to y_0 . Since we can transpose the pair (x_0, y_0) with C_0 we can

also transpose the pair (a_0, b_0) with G_0 . This clearly implies that we can transpose the pair (a_t, b_t) with G .

In order to conclude, it suffices to prove that, in the same situation as above, given any pair (c_t, d_t) of points in Γ_t , there is a permutation in G which sends the pair (c_t, d_t) to a pair (a_t, b_t) such that the points of its limiting pair (a_0, b_0) both sit on C_n^2 . We prove that acting with G we can move c_t to a point a_t such that its limit a_0 sits in C_n^2 . The same proof, applied to the stabilizer of a_t , will show that we can also move d_t to b_t as above.

Suppose c_t tends to a point c_0 on a line C_j^i , say on C_1^1 . The lines C_1^1 and C_1^2 meet at the point q_1 and, locally around q_1 , we have an open, smooth neighborhood A_t of C_t which tends to the union of two neighborhoods of q_1 on C_1^1 and C_1^2 respectively, thus acquiring a node at q_1 . By moving π_t on S_t^* close to π_0 , we can move the point c_t describing the whole open subset A_t of C_t . Therefore, by moving π_t in this way, we can let the limit point c_0 of c_t land on the line C_1^2 instead of C_1^1 . Our proof is thus finished.

Problem 2.3. Theorem 2.2 suggests the following general problem, which would be interesting to investigate. Let S be a smooth surface of degree d in $\mathbf{P}^3$ and let C be an irreducible curve of degree m on it. Under which conditions is the group $G(C, S)$ equal to S_m ? Notice that $G(C, S)$ is certainly not equal to S_m if S is a smooth quadric. In this case the section of any curve C on S with a tangent plane to S is formed by two sets of collinear points and the monodromy does not exchange points of the two sets. A similar situation takes place on any ruled surface S of degree d , but notice that then S is singular as soon as $d \geq 3$.

3. EXISTENCE OF REGULAR COMPONENTS

Our objective in this section is to prove our main result:

Theorem 3.1. *Let S be a general surface in $\mathbf{P}_d$, $d \geq 4$. For all $n \geq d$ and $\delta \leq N(d, n)$ there is an irreducible regular component of $V_{n, \delta}(S)$.*

As we know from §1, it suffices to prove the theorem for $\delta = N(d, n)$, which means that we have to exhibit an irreducible curve C in $|\mathcal{O}_S(n)|$ with $N(d, n)$ nodes and no other singularities, and with the nodes imposing independent conditions to $|\mathcal{O}_S(n)|$. Such a curve C gives rise to an isolated, reduced point of $V_{n, N(d, n)}^0$. We will say that such a curve C is *isolated*. We will start from Chen's theorem and prove the existence of these isolated curves inductively. Actually, we will inductively prove the following statement:

Theorem 3.2. *Let S be a general surface in $\mathbf{P}_d$, $d \geq 4$. For all $n \geq d$ and there are isolated curves C in $V_{n, N(d, n)}^0$ such that $G(C, S)$ is the full symmetric group S_{nd} .*

Our strategy for the proof of theorem 3.2 is the following. We consider a degenerate surface S_0 in $\mathbf{P}_d$ given as $S_0 = F \cup \pi$ where $F \in \mathbf{P}_{d-1}$ and π is a plane. We inductively require the existence of a surface $G \in \mathbf{P}_n$ such that $C_0^1 = F \cap G$ is irreducible, nodal and isolated with $N(d-1, n)$ nodes. Furthermore we want $C_0^2 = \pi \cap G$ nodal, with $(n-d+2)(n-d+3)/2$ nodes. Moreover we need that C_0^1 and C_0^2 intersect transversally at $n(d-1)$ points of $F \cap \pi$. Then we consider the curve $C = C_0^1 \cup C_0^2 \subset S_0 \subset G$ and we see that C deforms on a general surface

S of degree d to an irreducible, $N(d, n)$ -nodal curve C . Using induction we can prove that C is isolated and that $G(C, S)$ is the full symmetric group. However the main point is to start by constructing a *suitably good* surface S_0 . Let us see what the construction of S_0 amounts to.

First of all we make a remark. If theorem 3.2 holds, then there is a not empty open subset U of $\mathbf{P}_d$ over which the isolated curves of the statement fit together, giving at least one irreducible component V_d of the universal Severi variety $\mathcal{V}_{n, N(d, n)}^0$, which dominates $\mathbf{P}_d$ with 0-dimensional general fibers.

We need some more notation.

Fix numbers n, d with $n + 1 \geq d \geq 5$. Assume that theorem 3.2 holds for $d - 1$ and fix a component V_{d-1} of the universal Severi variety $\mathcal{V}_{n, N(d-1, n)}^0$, dominating $\mathbf{P}_{d-1}$ with finite fiber over the general point of $\mathbf{P}_{d-1}$.

Identifying a point $G \in \mathbf{P}_n$ with the surface in $\mathbf{P}^3$ parametrized by G , we let:

$\bar{X} \subset \mathbf{P}_n \times \mathbf{P}_{d-1}$ be a component of the variety of pairs (G, F) such that $G \cap F$ is an irreducible nodal curve in the fiber over F of the component V_{d-1} of the universal Severi variety;

X be the projection of $\bar{X}$ to $\mathbf{P}_n$;

$\bar{Y} \subset \mathbf{P}_n \times (\mathbf{P}^3)^*$ be the variety of pairs (G, π) with $G \cap \pi$ irreducible nodal plane curve with $(n - d + 2)(n - d + 3)/2$ nodes;

Y be projection of $\bar{Y}$ to $\mathbf{P}_n$;

$W \subset \mathbf{P}_n$ be the variety which parametrizes all reducible surfaces of degree n which split into a union $F \cup Q$ where F has degree $d - 1$, Q has degree $n - d + 1$ and both are irreducible;

$\tilde{\mathbf{P}}$ be the blow-up of $\mathbf{P}_n$ along the closure of W ;

E be the exceptional divisor of this blow-up;

$\tilde{X}, \tilde{Y}$ be the strict transforms of X, Y ;

$\tilde{W}$ be $\tilde{X} \cap E$.

As indicated in the rough sketch of proof of theorem 3.2 we gave above, the surface S_0 we need corresponds, with our notation, to a point of $X \cap Y$. In order to prove the existence of *suitably good* such surfaces, we will use degeneration to further reducible surfaces, and we will actually work on $\tilde{\mathbf{P}}$.

First we need a few preliminary results.

Proposition 3.3. (a) W is irreducible, of dimension

$$\dim W = \dim \mathbf{P}_{d-1} + \dim \mathbf{P}_{n-d+1} =$$

$$= 2 + \frac{1}{6}(n^3 - 3n^2d + 3nd^2 + 9n^2 + 12d^2 - 18nd + 26n - 24d).$$

(b) The tangent space to W at a general point $F \cup Q$ is naturally isomorphic to $H^0 \mathcal{I}_\gamma(n)/(F \cup Q)$ where γ is the curve $F \cap Q$.

(c) If theorem 3.2 holds for $d - 1$ then:

$$\dim \tilde{W} = \dim W + 1$$

Proof. (a) is trivial. (b) see §2 of [CHM]: a particular case is treated there, but the argument runs in general in the same way.

In order to compute the dimension of $\bar{X}$, project it to $\mathbf{P}_{d-1}$ and look at the fibers. On $F \in \mathbf{P}_{d-1}$ general, there are only finitely many nodal curves in $|\mathcal{O}_F(n)|$ with $N(d-1, n)$ nodes, so the fiber over F is a finite union of projective spaces $\mathbf{P}_{n-d+1}$. Since $\bar{X}$ dominates $\mathbf{P}_{d-1}$, the assertion clearly follows.

Proposition 3.4. *Y is irreducible, of codimension*

$$\text{codim}_{\mathbf{P}_n}(Y) = \frac{(n-d)(n-d+5)}{2}.$$

Proof. By [H], on a general plane π , the Severi variety $V := V(n, (n-d+2)(n-d+3)/2)$ is irreducible of codimension $(n-d+2)(n-d+3)/2$ in $|\mathcal{O}_\pi(n)|$. $\bar{Y}$ dominates V , with general fibers which are open subsets of projective spaces of fixed dimension, hence $\bar{Y}$ is irreducible and its dimension can be easily computed.

By the assumptions, we have $(n-d+2)(n-d+3) \geq 3$ and the general element of V is a nodal plane curve whose nodes are not aligned. Then the projection $\bar{Y} \rightarrow Y$ has finite fibers. In fact given a smooth surface G of degree $n \geq 2$ and given a nodal plane section C of G with at least three nodes which are not aligned, then clearly C is an isolated curve of the corresponding Severi variety.

Proposition 3.5. *Assume theorem 3.2 holds for $d-1$. Then W is a proper subvariety of codimension 1 in the closure of X .*

Proof. Consider a general element $F \cup Q \in W$. Since theorem 3.2 holds for $d-1$, there is some $G \in X$ such that $(F, G) \in \bar{X}$. Taking F general, we may assume that also G is general in X .

Consider a general element G' of the linear system spanned by $F \cup Q$ and G . Since $G \cap F = G' \cap F$ and G is general in X , then (F, G') belongs to $\bar{X}$, so that $G' \in X$. Thus in the limit, we get $F \cup Q \in X$. This proves that W is a proper subvariety of the closure of X .

Now, by proposition 3.3, (c), we have:

$$\dim W < \dim X \leq \dim \bar{X} = \dim W + 1$$

which concludes the proof of the proposition.

Remark 3.6. With a similar argument, we may show that a general element of X is smooth. Indeed, if $(F, G) \in \bar{X}$ is general, just as observed above, a general element in the linear system generated by G and by surfaces of type $F \cup Q$ must belong to X . The assertion easily follows by applying Bertini's theorem.

Proposition 3.7. *For $F \cup Q \in W$ general, the fiber of the exceptional divisor E over $F \cup Q$ is canonically isomorphic to $\mathbf{P}(H^0 \mathcal{O}_\gamma(n))$, where γ is the curve $F \cap Q$.*

Proof. The proof runs as in proposition 1 of [CHM] where a particular case is treated.

Proposition 3.8. *Suppose theorem 3.2 holds for $d-1$. Then the general fiber of the projection $\tilde{W} \rightarrow W$ is finite.*

Proof. Since W belongs to the closure of X , the projection $\tilde{W} \rightarrow W$ is dominant. By proposition 3.2

$$\dim W = \dim X - 1 = \dim \tilde{W}$$

because X is not contained in the exceptional divisor.

Remark 3.9. We can say more about the fibers of the projection $\tilde{W} \rightarrow W$.

Take general $F \in \mathbf{P}_{d-1}$, $Q \in \mathbf{P}_{n-d+1}$ and take $G \in \mathbf{P}_n$ such that (F, G) is a general point of $\tilde{X}$. The pencil spanned by $F \cup Q$ and G determines, as we saw in proposition 3.5, a tangent vector to X which is also a normal vector to W in $\mathbf{P}_n$, by proposition 3.3 (b). It follows that this pencil determines an element ψ of the fiber of $\tilde{W}$ over $F \cup Q$.

Now consider $\gamma = F \cap Q$ and think of ψ as an element of the fiber of E over $F \cup Q$, which is identified with $\mathbf{P}(H^0 \mathcal{O}_\gamma(n))$. By construction, ψ corresponds to the divisor $\gamma \cap G$ on γ . In other words, for F general, divisors in $|\mathcal{O}_\gamma(n)|$, corresponding to elements of $\tilde{W}$ are obtained by intersecting γ with the curves in $|\mathcal{O}_F(n)|$ having $N(d-1, n)$ nodes.

Lemma 3.10. [Hi] *Let $\mathcal{C}$ be a flat family of curves, whose general element is rational and irreducible. Let C_0 be a special, possibly reducible, member of the family. Then every irreducible component of C_0 is rational.*

Lemma 3.11. *Let F be general in $\mathbf{P}_{d-1}$, $d > 4$, and let $C \in |\mathcal{O}_F(n)|$ be an isolated nodal curve with $N(d-1, n)$ nodes, such that $G(C, S)$ is the full symmetric group.*

Take a general plane π tangent to F and take a set Z of $3(n-d+1) - 1$ points of $\Gamma = C \cap \pi$. Then there exists a finite, positive number of rational nodal curves of degree $n-d+1$ on π , with nodes outside F , passing through Z . More precisely, for each one of these rational nodal curves the union of the set of nodes and of the set Z gives independent conditions to the curves of degree $n-d+1$.

Proof. The family of rational nodal curves of degree $n-d+1$ in π is irreducible of dimension $3(n-d+1) - 1$. Thus, if we fix a set Z of $3(n-d+1) - 1$ points in Γ , we do get an element, in the closure of this family, which passes through Z . We need such an element to be nodal, irreducible, isolated, with its nodes off F .

Let $\mathcal{C}$ be any family of curves on π which is *rationally determined*, i.e. it is defined over the field of rational functions of π . The condition on $G(C, S)$ implies that Γ is in uniform position with respect to any rationally determined family $\mathcal{C}$ of curves on π , hence Z and Γ impose the same number of conditions to $\mathcal{C}$, unless Z imposes independent conditions to $\mathcal{C}$. By Hironaka's lemma 3.2, a degeneration of rational curves cannot contain the curve $F \cap \pi$, since we may assume that $F \cap \pi$ has positive genus. It follows that Z must impose independent conditions to the family $\mathcal{C}$ of rational curves in $|\mathcal{O}_\pi(n-d+1)|$ and to every subfamily of $\mathcal{C}$ which is rationally determined on π .

This implies that:

- (i) Z is contained in a finite number of rational curves $D \in |\mathcal{O}_\pi(n-d+1)|$;
- (ii) D cannot have singularities worse than nodes, since curves with such singularities fill a proper, closed subfamily of $\mathcal{C}$;
- (iii) similarly, we get that the set Z' of nodes of D does not intersect $F \cap \pi$;
- (iv) $Z \cup Z'$ impose independent conditions to curves of degree $n-d+1$.

We are now in a position to prove the existence of our surface $S = E \cup \pi$.

Lemma 3.12. *Assume theorem 3.2 holds for $d - 1$. Then there is an irreducible component B of $\tilde{Y} \cap \tilde{X}$ not contained in E , whose general point G , which we may identify with its image in $\mathbf{P}_n$, has the following properties:*

- (i) G is irreducible;
 - (ii) there is a plane π such that $\pi \cap G$ is irreducible, nodal, with $(n-d+2)(n-d+3)/2$ nodes which impose independent conditions to the curves of degree $n-d+1$, i.e. there are no such curves through these nodes;
 - (iii) there is a surface $F \in \mathbf{P}_{d-1}$ with $F \cap G$ irreducible, nodal and rigid, with $N(d-1, n)$ nodes;
 - (iv) the nodes of $F \cap G$ and $\pi \cap G$ are distinct and G is transversal to $F \cap \pi$.
- Furthermore, by moving G in B , we may assume F general in $\mathbf{P}_{d-1}$.

Proof. Look at $\tilde{W} \cap \tilde{Y}$. This intersection has a component A whose general element is $F \cup Q$, obtained by taking generically:

- a surface $F \in \mathbf{P}_{d-1}$;
- a plane π tangent to F ;
- a nodal curve $C = F \cap G'$ for some G' with $(F, G') \in \tilde{X}$;
- a subset $Z \subset \Gamma = C \cap \pi$ of degree $3(n-d+1) - 1$;
- a nodal rational curve $D \in |\mathcal{O}_\pi(n-d+1)|$, containing Z , with nodes off $F \cap \pi$ (see lemma 3.11);
- a surface Q of degree $n-d+1$, such that $Q \cap \pi = D$.

In fact in this way we found reducible surfaces $F \cup Q$ with a $N(d-1, n)$ -tangent surface G' of degree n , which defines the element of $\tilde{W}$ associated to $\psi = G' \cap C$ (see remark 3.8). Notice that π is tangent to F , tangent to Q at $(n-d)(n-d-1)/2$ distinct points and passes through a subset Z formed by $3(n-d+1) - 1$ points of ψ . Then, by the results of [Ch] and [R], the curve $\pi \cap (F \cup Q)$ is the limit of a nodal plane curve D with

$$1 + \frac{(n-d)(n-d-1)}{2} + 3(n-d+1) - 1 = \frac{(n-d+2)(n-d+3)}{2}$$

nodes lying on some irreducible surface G of degree n which degenerates to $F \cup Q$.

Let us compute the dimension of A . We have $\dim \mathbf{P}_{d-1}$ parameters for the choice of F , two parameters for the choice of π , no parameters for the choice of C , Z and D and $\dim \mathbf{P}_{n-d}$ parameters for the choice of the surface Q passing through the plane curve D . Thus:

$$\begin{aligned} \dim A &= 2 + \frac{1}{6}(n^3 - 3n^2d + 3nd^2 + 6n^2 - 9nd + 9d^2 + 11n - 9d) = \\ &= \dim W - \operatorname{codim} Y = \dim \tilde{X} + \dim \tilde{Y} - \dim \mathbf{P}_n - 1. \end{aligned}$$

It follows that there is a component B of $\tilde{X} \cap \tilde{Y}$ which contains A and is not contained in E . The general point G of B must be irreducible, since the elements of A split in two irreducible components and G cannot split in the same way. The plane π and the surface F exist and fulfill the transversality conditions (iv), since this happens for general points in A .

Finally observe that, by lemma 3.10, there are no curves of degree $n-d+1$ on π , other than $\pi \cap Q$, passing through Z and through the set nodes of $\pi \cap Q$. Since $\pi \cap Q$ misses the point where π is tangent to F , the nodes of the curve $\pi \cap (F \cup Q)$

- [CH] L. Caporaso, J. Harris, Degrees of Severi varieties, pre-print, 1996 (Alg. Geom. 9608025)
- [Chen] Xi Chen, Rational curves on K3 surfaces, Thesis, Harvard University, 1997
- [CS] L. Chiantini, E. Sernesi, Nodal curves on surfaces of general type, Math. Ann., 307 (1997), 41-56
- [CHM] C. Ciliberto, J. Harris, R. Miranda, General components of the Noether-Lefschetz locus and their density in the space of all surfaces, Math. Ann., 282 (1988), 667-680
- [GLS] G.M. Greuel, Ch. Lossen, E. Shustin, Geometry of families of nodal curves on the blown-up projective plane, pre-print, 1997 (Alg. Geom. 9704010)
- [H] J. Harris, On the Severi problem, Inventiones Math., 84 (1986), 445-461
- [Hi] H. Hironaka, On the arithmetic genera and the effective genera of algebraic curves, Mem. College Sci. Univ. Kyoto (A) 30 (1957), 177-195
- [K] M. Kontsevich, Enumeration of rational curves via torus action, pre-print
- [R] Z. Ran, Enumerative geometry of singular plane curves, Inventiones Math. 97 (1989), 447-465
- [S] E. Sernesi, On the existence of certain families of curves, Inventiones Math. 75 (1984), 25-57
- [Se] F. Severi, Vorlesungen ueber algebraische Geometrie, Teubner, Leipzig, 1921
- [T1] A. Tannenbaum, Families of algebraic curves with nodes, Compositio Math. 41 (1980), 107-126
- [T2] A. Tannenbaum, Families of curves with nodes on K3 surfaces, Math. Ann. 260 (1982), 239-253
- [YZ] S-T Yau, E. Zaslow, BPS states, string duality, and nodal curves on K3, pre-print, 1996